



Bayer CropScience LP  
 P.O. Box 12014, 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)

MOVENTO™

For Use On Bulb Onion For Management of Onion Thrips

SECTION 18 EMERGENCY EXEMPTION

File Symbol: 09-ID-06

## FOR DISTRIBUTION AND USE ONLY IN THE STATE OF IDAHO UNDER SECTION 18 EMERGENCY EXEMPTION

These use directions must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for MOVENTO™ (Registration Number 264-1050).

Any adverse effects resulting from the use of MOVENTO™ under this specific exemption must be immediately reported to the Idaho Department of Agriculture.

### EFFECTIVE PERIOD:

This specific exemption becomes effective 05/29/09 and expires on 09/15/09

A maximum of 9,000 acres of dry bulb onions may be treated in Idaho.

### DIRECTIONS FOR USE

CROP	PEST	Rate of MOVENTO™ fluid ounces/Acre
Onion (bulb)	Onion thrips	5.0

#### Notes and Restrictions

Movento may be applied by ground, air, or chemigation as directed on the Movento Section 3 label.

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum MOVENTO allowed per crop season: **10 fluid ounces/Acre (0.16 lb ai/A)**

No more than two applications are permitted per acre per season.

Onions must be harvested prior to bloom.

To minimize drift to adjacent fields, use ground applications whenever possible.

To the extent possible, local beekeepers should be notified prior to applying this chemical.

Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Apply specified dosage of MOVENTO as needed for control. Treatment must be made when thrips populations begin to build and before a damaging population becomes established. Best results will be obtained when two applications are made on a 7-10 day interval, incorporated into a seasonal program where effective products with different modes of action are utilized in rotational sequences for management of thrips populations. **MOVENTO must be tank-mixed with a spray adjuvant having spreading and penetrating properties to maximize leaf uptake and systemicity of the active ingredient within treated plants; please contact your local Bayer CropScience representative or PCA for specific recommendations.**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*